



Client: Newberg Veterinary Hospital	Patient: Giorgio
Email: frontdesk@newbergvet.com	Species/Breed: Dog / Labradoodle
Phone/Fax: 5035388303 / 503-538-8305	Age/Gender: 10 Month(s) / male
Owner: Dana Logan	Submitted Date: 11/07/2019 04:55:10 PM
Practitioner: Mark Weber	Completed Date: 11/08/2019 04:32:54 PM

Case Status/Priority: Completed / Regular

Case History:

hip and elbow evaluation

Case Report:

471259 KW/rb
Giorgio Logan

Six digitalized radiographic images containing projections of the pelvis and elbows were submitted for evaluation. The radiographs are dated 11/7/19.

Radiographic Findings:

Both femoral heads well seated in their respective acetabula with greater than 50% of the femoral heads located medial to the dorsal acetabular rims.

Both femoral heads are congruous with the acetabula; however, there is slight widening of the medial aspect of both coxofemoral joints.

No osteophyte formation is present on the femoral necks or acetabula.

Alignment of the patellae with the femurs is normal.

Muscle volume in both hind legs is within normal limits.

The anconeal process of both ulnae and proximocranial aspect of both radii are smooth with no evidence of osteophyte formation.

No sclerosis is associated with the ulnar notches.

Both medial coronoid processes are smooth in contour and a fragment associated with the medial coronoid processes is not identified.

Diagnosis: Normal pelvis and elbows.

Comments: Radiographic changes to suggest elbow or hip dysplasia are not present. Based on the OFA scheme of evaluating hips, the hips would be rated as good in conformation. Radiographs obtained at two years of age will need to be evaluated to completely exclude elbow and hip dysplasia and to establish a final rating on the coxofemoral joints.

Consulting Radiologist:

Kent Wallace, DVM, Ms, DACVR

kwallace@twcny.rr.com



Client: Newberg Veterinary Hospital	Patient: Charlie
Email: frontdesk@newbergvet.com	Species/Breed: Dog / Labradoodle
Phone/Fax: 5035388303 / 503-538-8305	Age/Gender: 5 Month(s) / female
Owner: Dana Logan	Submitted Date: 11/07/2019 04:57:00 PM
Practitioner: Mark Weber	Completed Date: 11/08/2019 04:33:26 PM

Case Status/Priority: Completed / Regular

Case History:

hip and elbow evaluation

Case Report:

471260 KW/rb
Charlie Logan

Seven digitalized radiographic images containing projections of the pelvis and elbows were submitted for evaluation. The radiographs are dated 11/7/19.

Radiographic Findings:

Both femoral heads are located in their respective acetabula and are congruous with the acetabula rims. There is slight widening of the medial aspect of both coxofemoral joints with this change slightly more dramatic involving the left coxofemoral joint. Alignment of the patellae with the femurs is normal. Muscle volume in both hind legs is within normal limits. The anconeal process of both ulnae and proximocranial aspect of both radii are smooth with no evidence of osteophyte formation. No sclerosis is associated with the ulnar notches. Both medial coronoid processes are smooth in contour and a fragment associated with the medial coronoid processes is not identified.

Diagnosis: Normal pelvis and elbows.

Comments: Radiographic changes to suggest elbow or hip dysplasia are not present. Based on the OFA scheme of evaluating hips, the hips would be rated as good in conformation. Radiographs obtained at two years of age will need to be evaluated to completely exclude elbow and hip dysplasia and to establish a final rating on the coxofemoral joints.

Consulting Radiologist:

Kent Wallace, DVM, Ms, DACVR
kwallace@twcny.rr.com



Client: Newberg Veterinary Hospital
Email: frontdesk@newbergvet.com
Phone/Fax: 5035388303 / 503-538-8305
Owner: Dana Logan
Practitioner: Mark Weber

Patient: Amelia
Species/Breed: Dog / Golden Retriever
Age/Gender: 2 Year(s) / female
Submitted Date: 11/07/2019 04:58:20 PM
Completed Date: 11/08/2019 04:39:41 PM

Case Status/Priority: Completed / Regular

Case History:

hip and elbow evaluation

Case Report:

471261 KW/rb
Amelia Logan

Five digitalized radiographic images containing projections of the pelvis and elbows were submitted for evaluation. The radiographs are dated 11/7/19.

Radiographic Findings:

Both femoral heads are well seated in their respective acetabula with greater than 50% of the left femoral head and slightly greater than 50% of the right femoral head are located medial to the dorsal acetabular rims.

Both femoral heads are congruous with the acetabular rims; however, there is slight widening of the medial aspect of both coxofemoral joints.

No osteophyte formation is present on the femoral necks or acetabula.

Alignment of the patellae with the femurs is normal.

Muscle volume in both hind legs is within normal limits.

The anconeal process of both ulnae and proximocranial aspect of both radii are smooth with no evidence of osteophyte formation.

No sclerosis is associated with the ulnar notches.

Both medial coronoid processes are smooth in contour and a fragment associated with the medial coronoid processes is not identified.

Diagnosis: Normal pelvis and elbows.

Comments: Radiographic changes to suggest elbow or hip dysplasia are not present. Based on the OFA scheme of evaluating hips, the hips would be rated as good in conformation. The clinical status of the patient will determine if further assessments radiographically would be indicated to evaluate the pelvis and elbows.

Consulting Radiologist:

Kent Wallace, DVM, Ms, DACVR

kwallace@twcny.rr.com