



Client: Newberg Veterinary Hospital
Email: frontdesk@newbergvet.com
Phone/Fax: 5035388303 / 503-538-8305
Owner: Dana Logan
Practitioner: Mark Weber

Patient: Oliver
Species/Breed: Dog / Australian Labardoodle
Age/Gender: 12 Month(s) / male
Submitted Date: 02/07/2020 12:03:02 PM
Completed Date: 02/27/2020 03:46:24 PM

Case Status/Priority: Completed / Regular

Case History:

Hip Evaluation

Case Report:

479756 KW/ss

Seven digitalized radiographic images containing projections of the pelvis and elbows were submitted for evaluation. The radiographs are dated 2/7/20.

Radiographic Findings:

Both femoral heads are located in their respective acetabula with greater than 50% of the femoral heads located medially to dorsal acetabular rims.

Obvious incongruity between the femoral heads and acetabula is not identified; however, there is slight widening of the medial aspect of both coxofemoral joints.

No osteophyte formation is present on the femoral necks or acetabula.

Alignment of the patellae with the femurs is normal.

Muscle volume in both hind legs is within normal limits.

The anconeal process of both ulnae and proximocranial aspect of both radii are smooth with no evidence of osteophyte formation.

No sclerosis is associated with the ulnar notches.

Both medial coronoid processes are smooth in contour and a fragment associated with the medial coronoid processes is not identified.

Diagnosis: Normal pelvis and elbows.

Comments: Radiographic changes to suggest elbow or hip dysplasia are not present. Based on the OFA scheme of evaluating hips, the hips would be rated as good in conformation. Radiographs obtained at 2 years of age will need to be evaluated to completely exclude elbow and hip dysplasia and to establish a final rating on the coxofemoral joints.

Consulting Radiologist:

Kent Wallace, DVM, Ms, DACVR

kwallace@twcny.rr.com